



AYSO KANEOHE REGION 113

PLAYER INFORMATION FORM

N / R Player Name: _____ Parent Name: _____

Birthdate: _____ (BC) Gender: M / F Age Division: _____

Parent Phone: _____ Parent email: _____

General Info/Check-In Station: _____ (please check when parent is given packet)

Uniform Station (circle size):

YXXS YXS YS YM YL AS AM AL AXL A2XL

Parent initials: _____ *

**When initialing, you are confirming uniform size*

Volunteer Station:

We are an all-volunteer organization and without volunteers we would not be able to run a soccer program in the Kaneohe area. Any position you can assist in is greatly appreciated!

Please circle:

COACH: _____ (AGE DIVISION/GENDER)

TEAM PARENT

ASSISTANT COACH: _____ (AGE DIVISION/GENDER)

REFEREE


BOARD MEMBER: Assist in running of the region. Various positions available:

- 8U AGC (age group coordinator): This position is responsible for coordinating 8U division
- MARKETING COORDINATOR: The position is responsible for promoting Kaneohe AYSO in our community.

PICTURE DAY HELPER: Assist with team picture taking day(s). Normally held over 2 weekends in August/Sept

SPECIAL EVENTS HELPER: Assist with various Region events such as Player's Clinic, Soccerfest or Kaneohe Christmas Parade

Parent/Athlete Concussion Information Sheet



A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by bump, blow or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports **one or more** symptoms of concussion listed

Did You Know?

- Most concussions occur *without* loss of consciousness
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

below after a bump, blow, or jolt to the head or body, s/he should be kept out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

SIGNS OBSERVED BY COACHING STAFF

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (*even briefly*)
- Shows mood, behavior, or personality changes
- Can't recall events *prior* to hit or fall
- Can't recall events *after* hit or fall

SYMPTOMS REPORTED BY ATHLETES

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or "feeling down"

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (*even a brief loss of consciousness should be taken seriously*)

Remember

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

It's better to miss one game than the whole season. For more information on concussions, visit: www.cdc.gov/Concussion

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. *They can even be fatal.*

Student-Athlete Name Printed

Student-Athlete Signature

Date

Parent or Legal Guardian Printed

Parent or Legal Guardian Signature

Date